



PHOX2B gene

paired like homeobox 2b

Normal Function

The *PHOX2B* gene provides instructions for making a protein that acts early in development to help promote the formation of nerve cells (neurons) and regulate the process by which the neurons mature to carry out specific functions (differentiation). The protein is active in the neural crest, which is a group of cells in the early embryo that give rise to many tissues and organs. Neural crest cells migrate to form parts of the autonomic nervous system, which controls body functions such as breathing, blood pressure, heart rate, and digestion. Neural crest cells also give rise to many tissues in the face and skull, and other tissue and cell types.

The protein produced from the *PHOX2B* gene contains two areas where a protein building block (amino acid) called alanine is repeated multiple times. These stretches of alanines are known as polyalanine tracts or poly(A) tracts.

Health Conditions Related to Genetic Changes

congenital central hypoventilation syndrome

Most *PHOX2B* gene mutations that cause congenital central hypoventilation syndrome (CCHS) add extra alanines to the polyalanine tracts in the PHOX2B protein. This type of mutation is called a polyalanine repeat expansion. Other types of *PHOX2B* gene mutations have been identified in 8 to 10 percent of individuals with this disorder.

Mutations are believed to interfere with the PHOX2B protein's role in promoting neuron formation and differentiation, especially in the autonomic nervous system, resulting in the problems regulating breathing and other autonomic nervous system dysfunction seen in CCHS.

neuroblastoma

Several mutations in the *PHOX2B* gene have been identified in people with neuroblastoma, a type of cancerous tumor composed of immature nerve cells (neuroblasts). Neuroblastoma and other cancers occur when a buildup of genetic mutations in critical genes—those that control cell proliferation or differentiation—allow cells to grow and divide uncontrollably to form a tumor. In most cases, these genetic changes are acquired during a person's lifetime, called somatic mutations. Somatic mutations are present only in certain cells and are not inherited. Less commonly, gene mutations that increase the risk of developing cancer

can be inherited from a parent. Both types of mutation occur in neuroblastoma. Somatic mutations in the *PHOX2B* gene increase the risk of developing sporadic neuroblastoma, and inherited mutations in the *PHOX2B* gene increase the risk of developing familial neuroblastoma.

In some people with neuroblastoma, mutations in the *PHOX2B* gene change a single protein building block (amino acid) in the PHOX2B protein. Other affected individuals may have an addition or deletion of several DNA building blocks (nucleotides) in the *PHOX2B* gene. Addition or deletion of nucleotides changes the sequence of amino acids in the PHOX2B protein. All of these types of mutations have been found in familial and sporadic neuroblastoma. The mutations are believed to interfere with the PHOX2B protein's role in promoting nerve cell differentiation, which results in an excess of immature nerve cells and leads to neuroblastoma.


other disorders

Variations in the *PHOX2B* gene have been associated with increased risk of certain other disorders involving the autonomic nervous system and tissues derived from the neural crest. Particular *PHOX2B* gene variations have been identified in people who have both congenital central hypoventilation syndrome and a dysfunction of the nerves in the intestine called Hirschsprung disease (this combination of disorders is often called Haddad syndrome). The nerve problems in Hirschsprung disease result in severe constipation, intestinal blockage, and enlargement of the colon in affected individuals. *PHOX2B* gene mutations have also been identified in people with both neuroblastoma and Hirschsprung disease. In addition, *PHOX2B* gene variations have been identified in some babies who died of sudden infant death syndrome (SIDS). *PHOX2B* gene variations likely affect the regulation of neuron differentiation in early development, resulting in an increased risk of these disorders.

Chromosomal Location

Cytogenetic Location: 4p13, which is the short (p) arm of chromosome 4 at position 13

Molecular Location: base pairs 41,744,082 to 41,748,970 on chromosome 4 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- NBLST2
- NBPhox
- neuroblastoma paired-type homeobox protein
- neuroblastoma Phox
- paired-like homeobox 2b
- paired mesoderm homeobox 2b
- Phox2b
- PHOX2B homeodomain protein
- PHX2B_HUMAN
- PMX2B

Additional Information & Resources

GeneReviews

- Congenital Central Hypoventilation Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1427>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28PHOX2B%5BTIAB%5D%29+OR+%28paired-like+homeobox+2b%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

OMIM

- HIRSCHSPRUNG DISEASE, SUSCEPTIBILITY TO, 1
<http://omim.org/entry/142623>
- PAIRED-LIKE HOMEODOMAIN PROTEIN 2B
<http://omim.org/entry/603851>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/PHOX2BID126ch4p13.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=PHOX2B%5Bgene%5D>

- HGNC Gene Family: PRD class homeoboxes and pseudogenes
<http://www.genenames.org/cgi-bin/genefamilies/set/521>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=9143
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/8929>
- UniProt
<http://www.uniprot.org/uniprot/Q99453>

Sources for This Summary

- Antic NA, Malow BA, Lange N, McEvoy RD, Olson AL, Turkington P, Windisch W, Samuels M, Stevens CA, Berry-Kravis EM, Weese-Mayer DE. PHOX2B mutation-confirmed congenital central hypoventilation syndrome: presentation in adulthood. *Am J Respir Crit Care Med*. 2006 Oct 15; 174(8):923-7. Epub 2006 Jul 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16873766>
- Berry-Kravis EM, Zhou L, Rand CM, Weese-Mayer DE. Congenital central hypoventilation syndrome: PHOX2B mutations and phenotype. *Am J Respir Crit Care Med*. 2006 Nov 15;174(10): 1139-44. Epub 2006 Aug 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16888290>
- Gaultier C, Trang H, Dauger S, Gallego J. Pediatric disorders with autonomic dysfunction: what role for PHOX2B? *Pediatr Res*. 2005 Jul;58(1):1-6. Epub 2005 May 18. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15901893>
- GeneReview: Congenital Central Hypoventilation Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1427>
- Gronli JO, Santucci BA, Leurgans SE, Berry-Kravis EM, Weese-Mayer DE. Congenital central hypoventilation syndrome: PHOX2B genotype determines risk for sudden death. *Pediatr Pulmonol*. 2008 Jan;43(1):77-86.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18041756>
- Miao X, Garcia-Barceló MM, So MT, Leon TY, Lau DK, Liu TT, Chan EK, Lan LC, Wong KK, Lui VC, Tam PK. Role of RET and PHOX2B gene polymorphisms in risk of Hirschsprung's disease in Chinese population. *Gut*. 2007 May;56(5):736.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17440194>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1942142/>
- Parodi S, Bachetti T, Lantieri F, Di Duca M, Santamaria G, Ottonello G, Matera I, Ravazzolo R, Ceccherini I. Parental origin and somatic mosaicism of PHOX2B mutations in Congenital Central Hypoventilation Syndrome. *Hum Mutat*. 2008 Jan;29(1):206.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18157832>
- Raabe EH, Laudenslager M, Winter C, Wasserman N, Cole K, LaQuaglia M, Maris DJ, Mosse YP, Maris JM. Prevalence and functional consequence of PHOX2B mutations in neuroblastoma. *Oncogene*. 2008 Jan 17;27(4):469-76. Epub 2007 Jul 16.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17637745>

- Reiff T, Tsarovina K, Majdazari A, Schmidt M, del Pino I, Rohrer H. Neuroblastoma phox2b variants stimulate proliferation and dedifferentiation of immature sympathetic neurons. *J Neurosci*. 2010 Jan 20;30(3):905-15. doi: 10.1523/JNEUROSCI.5368-09.2010.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20089899>
- Repetto GM, Corrales RJ, Abara SG, Zhou L, Berry-Kravis EM, Rand CM, Weese-Mayer DE. Later-onset congenital central hypoventilation syndrome due to a heterozygous 24-polyalanine repeat expansion mutation in the PHOX2B gene. *Acta Paediatr*. 2009 Jan;98(1):192-5. doi: 10.1111/j.1651-2227.2008.01039.x. Epub 2008 Sep 16.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18798833>
- Trochet D, Hong SJ, Lim JK, Brunet JF, Munnich A, Kim KS, Lyonnet S, Goridis C, Amiel J. Molecular consequences of PHOX2B missense, frameshift and alanine expansion mutations leading to autonomic dysfunction. *Hum Mol Genet*. 2005 Dec 1;14(23):3697-708. Epub 2005 Oct 25.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16249188>
- Weese-Mayer DE, Berry-Kravis EM, Ceccherini I, Rand CM. Congenital central hypoventilation syndrome (CCHS) and sudden infant death syndrome (SIDS): kindred disorders of autonomic regulation. *Respir Physiol Neurobiol*. 2008 Dec 10;164(1-2):38-48. doi: 10.1016/j.resp.2008.05.011. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18579454>
- Weese-Mayer DE, Berry-Kravis EM, Marazita ML. In pursuit (and discovery) of a genetic basis for congenital central hypoventilation syndrome. *Respir Physiol Neurobiol*. 2005 Nov 15;149(1-3):73-82. Epub 2005 Jul 28. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16054879>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/PHOX2B>

Reviewed: August 2012
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services